

## Short communication. The antifeedant activity of natural plant products towards the larvae of *Spodoptera littoralis*

M. C. Ballesta-Acosta<sup>1</sup>, M. J. Pascual-Villalobos<sup>1\*</sup> and B. Rodríguez<sup>2</sup>

<sup>1</sup> Instituto Murciano de Investigación y Desarrollo Agrario y Alimentario (IMIDA), Estación Sericícola.  
C/ Mayor, s/n. 30150 La Alberca (Murcia). Spain

<sup>2</sup> Instituto de Química Orgánica, CSIC. C/ Juan de la Cierva, 3. 28006 Madrid. Spain

### Abstract

Worms are becoming difficult pests to control since many insecticides are now forbidden and the present alternatives are insufficiently effective. Antifeedants of plant origin may have potential against such pests. In this work, several diterpenoids, flavones and coumarins extracted from plants were assayed as antifeedant agents against *Spodoptera littoralis* larvae. Among these compounds, four diterpenes (eriocephalin, salviacoccin, aethiopinone and oxocandesalvone) and four coumarins (oxypeucedanin, xanthotoxin, isoimperatorin and prangol) showed significant activity, whereas all the flavones tested were inactive. Some aspects of the structure-activity relationships of these compounds are briefly discussed.

**Additional key words:** coumarins, diterpenes, flavones, higher plants, secondary metabolites.

### Resumen

#### Comunicación corta. Actividad antialimentaria de fitoquímicos naturales en larvas de *Spodoptera littoralis*

En la actualidad, debido a que se están prohibiendo muchos insecticidas y a que las alternativas disponibles no son completamente eficaces, las orugas están resurgiendo como plagas difíciles de controlar. Los compuestos antialimentarios de origen vegetal se consideran con posibilidades para desarrollar nuevos agroquímicos. En este trabajo, diversos diterpenoides, flavonas y cumarinas procedentes de plantas han sido ensayados como antialimentarios frente a larvas de *Spodoptera littoralis*. Tres diterpenos (eriocefalina, salviacoccina, aetiopinona y oxocandesalvona) y cuatro cumarinas (oxipeucedanina, xantotoxina, isoimperatorina y prangol) presentaron una actividad significativa, mientras que las flavonas fueron inactivas. Se discuten brevemente algunos aspectos de la relación estructura-actividad.

**Palabras clave adicionales:** cumarinas, diterpenos, flavonas, metabolitos secundarios, plantas superiores.

Moths are common, polyphagous pests in Murcia, Andalusia and Valencia, the main vegetable producing regions of Spain. In these areas, leaf-feeding noctuid lepidopterans, including *Spodoptera exigua* Hübner, *Autographa gamma* L. and *Chrysodeixis chalcites* Esper., and borer worms such as *Helicoverpa armigera* Hübner, *Ostrinia nubilalis* Hübner (Lepidoptera, Pyralidae) and *Gorthina xanthenes*, are gradually causing more damage and are becoming hard to control since fewer chemicals are available and problems of resistance are emerging (Torres-Vila *et al.*, 2002).

The army worm, *Spodoptera littoralis* (Boisduval, 1833) (Lepidoptera: Noctuidae), is an important pest of cotton, tobacco and vegetable crops (tomato, pepper, lettuce, broccoli, cauliflower and artichoke). The larvae cause economically important damage since they feed on the leaves, fruits and buds from summer till the end of autumn in the field (or even year round when warm temperatures prevail during winter and spring) and year round in greenhouses (Rodríguez-Rodríguez *et al.*, 1996; Téllez *et al.*, 1996). Control methods include the use of agrochemicals such as pyrethroids, tebufen-

\* Corresponding author: [mjesus.pascual@carm.es](mailto:mjesus.pascual@carm.es)  
Received: 23-03-07; Accepted: 20-11-07.

Abbreviations used: AI (antifeedant indices); ppm (parts per million).

nocide and spinosad (organophosphates are gradually being banned), the use of *Bacillus thuringiensis* Berliner (Ramos-Gutiérrez et al., 2004), baculovirus (Baculoviridae) (Murillo et al., 2006), and finally the use of pheromones (under development).

Plants are important sources of natural products with the potential to be developed as commercial active ingredients for pest control. Antifeedants have been proposed as an alternative to synthetic insecticides (Belles et al., 1985). Clerodane diterpenoids are well known insect antifeedants isolated from plants of the genera *Ajuga*, *Leonororus*, *Salvia*, *Scutellaria*, *Stachys* and *Teucrium* (family Labiateae) (Merrit and Ley, 1992; De la Torre et al., 1994; Rodríguez-Hanh et al., 1994; Bremner et al., 1998). The most potent neoclerodanes are jodrelin and scutalpin, which are isolated from *Scutellaria* (Anderson et al., 1989). The coumarins have been much less studied although they too have been cited as insect antifeedants (Addor, 1994; Stevenson et al., 2003). In addition, a few papers document the antifeedant effects of flavones (Sharaby and Ammar, 1997) and other diterpenoids, for example those found in *Plectranthus* leaves (Wellsow et al., 2006).

This paper reports the antifeedant activity against *S. littoralis* of natural plant compounds belonging to different classes—clerodane, abietane, labdane and rosane diterpenoids, flavones and coumarins—with the aim of providing knowledge that may be of use in the development of new agrochemical products.

The phytochemicals used in this study are listed in Table 1 along with their plant sources.

*Spodoptera littoralis* was reared on a bean-based diet at  $26 \pm 1^\circ\text{C}$  under a 16L:8D photoperiod. Larvae were originally collected several years ago from an alfalfa field in Totana (Province of Murcia); since then, this population has been reared in the lab to supply insects for research purposes.

Experiments were performed in 9 cm diameter Petri dishes each containing a glass fibre disc (diameter 2.1 cm) (Whatman GF/A). One hundred microlitres of a 50 mM sucrose solution plus 100  $\mu\text{L}$  of a 100 ppm solution of the test compounds were applied to different discs (control discs received only the sucrose solution). A sixth instar larva was then added to each Petri dish. The discs (after the application of the products) were weighed before and after (18 h) the bioassay. Ten replicates were prepared for each compound assayed.

Following the methodology described by Simmonds et al. (1990), antifeedant indices (AI) were calculated according to the formula:

$$\text{AI} = [(C - T) / (C + T)] \times 100$$

where C = the weight (g) of the control disc consumed, and T = the weight (g) of the treated disc consumed. The AI ranged from -100 to +100. If a value of  $\text{AI} > 0$  was returned, the compound was deemed to be an antifeedant; if a result of  $\text{AI} < 0$  was returned, the compound was regarded as a phagostimulant. A mean AI was calculated for each assayed phytochemical. Comparisons between the eaten weight of the treated and control discs were made using the non-parametric Wilcoxon test.

The most active phytochemical was then assayed at different concentrations (4 doses, 2 replicates per dose, 100 larvae per replicate). Probit analysis (Robertson and Priesler, 1992) was then used to determine the doses with an antifeedant effect ( $\text{AI} \geq 50$ ) towards 10, 50 or 90% of the larvae.

Table 2 shows that the clerodane diterpenoids eriocephalin, salviacoccin and salvifarín, were active at 100 ppm, while the other compounds of this class had an AI value of around zero. The most active product ( $\text{AI} = 62.18$ ) was eriocephalin. Overall, the abietane diterpenoids had less of an antifeedant effect, although aethiopinone and 2-oxocandesalvone A returned statistically significant results ( $\text{AI} = 29.82$  and 23.02 respectively). In addition, larval feeding was reduced by the intake of 12-O-demethylcryptojaponol. As for the labdane and rosane diterpenoids tested, positive AI were obtained, although they were not much greater than those for the previously mentioned compounds (Table 2).

Eriocephalin, a compound isolated from *Teucrium eriocephalum* Wk, showed the greatest antifeedant effect. Table 3 shows the results of the probit analysis. The ED50 was 43 ppm, i.e., at this dose, an antifeedant effect (with  $\text{AI} \geq 50$ ) would be seen against 50% of larvae. European and Asian species of *Salvia* contain more abietane than clerodane diterpenoids (Rodríguez-Hanh et al., 1994). Aethiopinone is a strong anti-inflammatory substance (Hernández-Pérez et al., 1995) and possesses antimicrobial activity as well as cytotoxic activity towards cultured KB human carcinoma cells (Hernández-Pérez et al., 1999). Eriocephalin is known to be very active against *S. littoralis* (Simmonds et al., 1989; De la Torre et al., 1994). According to Geusken et al. (1983), substitutions in the decalin ring ( $4\alpha,18$ -oxirane) and a  $\beta$ -substituted furan at C-12 are key structural features in this activity.

In contrast to the results reported here for *S. littoralis*, Ortego et al. (1995) reported that teucvin (a 19-nor

**Table 1.** Phytochemicals tested

Class / compound	Plant source <sup>a</sup>	Reference
<i>Clerodane diterpenoids</i>		
Salviacoccin	<i>Salvia coccinea</i> Juss	Savona <i>et al.</i> (1982)
Salvifarín	<i>Salvia farinacea</i> Benth.	Eiguren <i>et al.</i> (1984)
Eriocéphalin	<i>Teucrium eriocéphalum</i> Wk.	Fayos <i>et al.</i> (1979)
Isoeriocephalin	<i>Teucrium lanigerum</i> Lag.	Fernández-Gadea <i>et al.</i> (1984)
Teuflin	<i>Teucrium chamaedrys</i> L.	Fernández-Gadea <i>et al.</i> (1983)
Teucvin	<i>Teucrium chamaedrys</i> L	Rodríguez <i>et al.</i> (1984a)
Teuflidin	<i>Teucrium chamaedrys</i> L.	Fernández-Gadea <i>et al.</i> (1983)
<i>Abietane diterpenoids</i>		
Royleanone	<i>Salvia phlomoides</i> Asso.	Hueso-Rodríguez <i>et al.</i> (1983)
7α-Acetoxyroyleanone	<i>Salvia phlomoides</i> Asso.	
Taxodione	<i>Salvia phlomoides</i> Asso.	
Taxodone	<i>Salvia phlomoides</i> Asso.	
Crytojaponol	<i>Salvia phlomoides</i> Asso.	
12-O-Demethylcryptojaponol	<i>Salvia phlomoides</i> Asso.	
Horminone	<i>Salvia pratensis</i> L.	Hueso-Rodríguez and Rodríguez (1989)
12-O-Methylcandelabrone	<i>Salvia candelabrum</i> Boiss.	Mendes <i>et al.</i> (1989)
12-O-Methylcandesalvone A	<i>Salvia candelabrum</i> Boiss.	
12-O-Methylcandesalvone B 4-methyl ester	<i>Salvia candelabrum</i> Boiss.	
Aethiopinone	<i>Salvia aethiopis</i> L.	Rodríguez <i>et al.</i> (1984b)
2-Oxocandesalvone A 12-O-methyl ether	<i>Salvia palaestina</i> L.	Hussein <i>et al.</i> (1997)
<i>Labdane diterpenoid</i>		
Hispanolone	<i>Ballota hirsuta</i> Benth.	Savona <i>et al.</i> (1978)
<i>Rosane diterpenoid</i>		
Lagascatriol	<i>Sideritis angustifolia</i> Lag.	Martín-Panizo <i>et al.</i> (1974)
<i>Flavones</i>		
Salvigenin	<i>Sideritis serrata</i> Lag.	Rodríguez and Martín-Panizo 1979
Eupatorin	<i>Teucrium pseudochamaepitys</i> L.	Savona <i>et al.</i> (1979)
Cirsilineol	<i>Sideritis mugronensis</i> Borja	Rodríguez (1977)
5-Hydroxy-6,7,3',4'-tetramethoxyflavone	<i>Sideritis mugronensis</i> Borja	
Gardenin D	<i>Sideritis mugronensis</i> Borja	
5,4'-Dihydroxy-6,7,8,3'-tetramethoxyflavone	<i>Sideritis mugronensis</i> Borja	
5-O-Demethylnobiletin	<i>Sideritis mugronensis</i> Borja	
<i>Coumarins</i>		
Isoimperatorin	<i>Cachrys trifida</i> Miller	Hernández and Rodríguez (1981)
Imperatorin	<i>Cachrys trifida</i> Miller	
Prantschimgin	<i>Cachrys trifida</i> Miller	
Bergapten	<i>Cachrys trifida</i> Miller	
Cnidilin	<i>Cachrys trifida</i> Miller	
Oxypeucedanin	<i>Cachrys trifida</i> Miller	
Xanthotoxin	<i>Cachrys trifida</i> Miller	
Prangol	<i>Cachrys trifida</i> Miller	

<sup>a</sup> All species are members of the family Labiatae, except for *Cachrys trifida* Miller which belongs to the family Umbelliferae.

**Table 2.** Antifeedant activity of phytochemicals (100 µL of a 100 ppm solution applied to 2.1 cm diam. discs) towards *Spodoptera littoralis*

Class / compound	AI (mean ± SEM) <sup>a</sup>	P <sup>b</sup>
<i>Clerodane diterpenoids</i>		
Salviacoccin	31.28 ± 17.669	0.0049*
Salvifarin	21.53 ± 19.179	0.1396
Eriocophilin	62.18 ± 11.338	0.0020*
Isoeriocephalin	0.39 ± 19.227	0.1797
Teuflin	4.14 ± 15.237	0.4043
Teucvin	0.25 ± 15.293	0.4219
Teuflidin	0.97 ± 22.587	0.1504
<i>Abietane diterpenoids</i>		
Royleanone	-37.20 ± 13.057	0.0420 <sup>#</sup>
7α-Acetoxyroyleanone	6.11 ± 22.509	0.2539
Taxodione	-25.50 ± 18.276	0.1396
Taxodone	16.43 ± 13.922	0.0654
Cryojaponol	-10.90 ± 10.591	0.0967
12-O-Demethylcryptojaponol	33.42 ± 14.813	0.0645
Horminone	-21.00 ± 16.222	0.0820
12-O-Methylcandelabrone	10.39 ± 11.255	0.2305
12-O-Methylcandesalvone A	5.04 ± 18.275	0.3262
12-O-Methylcandesalvone B 4-methyl ester	20.92 ± 14.096	0.1650
Aethiopinone	29.82 ± 11.861	0.0273*
2-Oxocandesalvone A 12-O-methyl ether	23.02 ± 19.113	0.0186*
<i>Labdane diterpenoid</i>		
Hispanolone	11.64 ± 10.338	0.1132
<i>Rosane diterpenoid</i>		
Lagascatriol	23.43 ± 20.134	0.4043
<i>Flavones</i>		
Salvigenin	-8.75 ± 13.404	0.5332
Eupatorin	-5.19 ± 19.096	0.5332
Cirsilineol	15.46 ± 17.058	0.3633
5-Hydroxy-6,7,3',4'-tetramethoxyflavone	-6.62 ± 15.876	0.4482
Gardenin D	-38.86 ± 24.109	0.0801
5,4'-Dihydroxy-6,7,8,3'-tetramethoxyflavone	38.31 ± 56.245	0.4043
5-O-Demethylnobiletin	-14.53 ± 15.156	0.3236
<i>Coumarins</i>		
Isoimperatorin	31.89 ± 22.921	0.0098*
Imperatorin	-31.38 ± 13.430	0.0420 <sup>#</sup>
Prantschimgin	-33.79 ± 19.592	0.0039 <sup>#</sup>
Bergapten	2.72 ± 17.996	0.4707
Cnidilin	8.86 ± 19.526	0.4707
Oxypeucedanin	41.92 ± 18.747	0.0137*
Xanthotoxin	43.99 ± 11.793	0.0186*
Prangol	26.78 ± 25.441	0.0195*

<sup>a</sup> Antifeedant index: AI = [(C - T) / (C + T)] × 100; C = consumed weight of control disc, T = consumed weight of treated disc; AI > 0 = antifeedant activity, A < 0 = phagostimulant activity. <sup>b</sup> Probability for C - T (Wilcoxon test). \* Statistically significant antifeedant activity (P < 0.05). <sup>#</sup> Statistically significant phagostimulant activity (P < 0.05).

**Table 3.** Regression analysis of eriocephalin antifeedant effect (AI ≥ 50) in *S. littoralis* larvae

<b>ED10<sup>a</sup> (ppm)</b>	<b>95% Confidence limits</b>	<b>ED50<sup>b</sup> (ppm)</b>	<b>95% Confidence limits</b>	<b>ED90<sup>c</sup> (ppm)</b>	<b>95% Confidence limits</b>
0.54	0.25-1.925	43	12-175	3469	1000- >10000

<sup>a,b,c</sup> Doses [ppm; solutions (100 µL) were applied to 2.1 cm diam. discs] with antifeedant effect (AI ≥ 50) towards 10%, 50% and 90% of larvae respectively.

neoclerodane) at 100 ppm caused a significant reduction in feeding in *Leptinotarsa decemlineata* larvae.

Salviacoccin and salvifarín have a saturated and α,β-unsaturated γ-lactone group respectively. According to Simmonds *et al.* (1996), this explains the activity of these compounds extracted from *Salvia*. These authors obtained an AI of 59 (at a dose of 100 ppm) for salvia-coccin – a stronger antifeedant effect than that seen in the present experiment (AI = 31.28).

The flavones had no significant effect on the feeding behaviour of the larvae (Table 2).

The coumarins were active against *S. littoralis*. Oxypeucedanin and xanthotoxin gave AI values of over 40 (Table 2). A remarkable antifeedant activity was recorded for isoimperatorin (AI = 31.89) and prangol (AI = 26.78). The coumarins are a class of naturally occurring antifeedants that have been less studied than the neoclerodanes. Calcagno *et al.* (2002) reported a deterrent effect of xanthotoxin on *S. littoralis*, with synergistic effects in mixtures with imperatorin.

When the structures of active and inactive coumarins, such as isoimperatorin and imperatorin or xanthotoxin and bergapten are compared, substitution groups at different positions (C-4 or C-9) account for the differences in antifeedant activity (Table 1).

Some papers report the medicinal potential of furocoumarins such as imperatorin, isoimperatorin, oxypeudanin and prantschimgin (Abad *et al.*, 2001; Hyuncheol *et al.*, 2002).

In summary, the active diterpenoids isolated from species of *Teucrium* and *Salvia* (Labiatae) and a number of coumarins from *Cachrys trifida* (Umbelliferae), appear to have antifeedant potential with respect to *S. littoralis*. More research is required in this area to help find alternatives to banned pesticides.

## Acknowledgments

The authors would like to thank the INIA for funding this work (project SC98-022) and the Fundación

Séneca (Murcia) for the doctoral fellowship awarded to M.C. Ballesta-Acosta.

## References

- ABAD M.J., DE LAS HERAS B., SILVAN A.M., PASCUAL R., BERMEJO P., RODRÍGUEZ B., VILLAR A.M., 2001. Effect of furocoumarins from *Cachrys trifida* on some macrophage functions. *J Pharm Pharmacol* 53, 1163-1168.
- ADDOR R.W., 1994. Insecticides. In: Agrochemicals from natural products (Godfrey C.R.A., ed). Marcel Dekker, Inc, NY. pp. 1-62.
- ANDERSON J.C., BLANEY W.M., COLE M.D., FELLOWS L.L., LEY S.V., SHEPPARD R.N., SIMMONDS M.S.J., 1989. The structure of two new clerodane diterpenoids potent insect antifeedants from *Scutellaria woronowii* (Juz): jodrellin A & B. *Tetrahedron Lett* 30, 4737-4740.
- BELLES X., CAMPS F., COLL J., PIULACHS M.D., 1985. Insect antifeedant activity of clerodane diterpenoids against larvae of *Spodoptera littoralis* (Boisd.) (Lepidoptera). *J Chem Ecol* 11, 1439-1445.
- BREMNER P., SIMMONDS M.S.J., BLANEY W.M., VEITCH N.C., 1998. Neo-clerodane diterpenoid insect antifeedants from *Ajuga reptans* cv. *Catkins Giant*. *Phytochemistry* 47, 1227-1232.
- CALCAGNO M.P., COLL J., LLORIA J., FAINI F., ALONSO-AMELOT M.E., 2002. Evaluation of synergism in the feeding deterrence of some furanocoumarins on *Spodoptera littoralis*. *J Chem Ecol* 28, 175-191.
- DE LA TORRE M.C., DOMÍNGUEZ G., RODRÍGUEZ B., PERALES A., SIMMONDS M.S.J., BLANEY W.M., 1994. Some remarkable reactions of the diterpene eriocephalin: neoclerodane derivatives with insect antifeedant activity. *Tetrahedron* 50, 13553-13566.
- EGUREN L., FAYOS J., PERALES A., SAVONA G., RODRÍGUEZ B., 1984. Salvifarín, X-ray structure determination of a *cis* neo-clerodane diterpenoid from *Salvia farinacea*. *Phytochemistry* 23, 466-467.
- FAYOS J., MARTÍNEZ-RIPOLL M., PATERNOSTRO M. P., PIOZZI F., RODRÍGUEZ B., SAVONA G., 1979. New clerodane diterpenoid from *Teucrium eriocephalum*. *J Org Chem* 44, 4992-4994.
- FERNÁNDEZ-GADEA F., PASCUAL C., RODRÍGUEZ B., SAVONA G., 1983. 6-Epitemucrin A, a neo-clerodane diterpenoid from *Teucrium chamaedrys*. *Phytochemistry* 22, 723-725.

- FERNÁNDEZ-GADEA F., RODRÍGUEZ B., SAVONA G., PIOZZI F., 1984. Isoerocephalin and 20-deacetyleriocephalin, neo-clerodane diterpenoids from *Teucrium lanigerum*. Phytochemistry 23, 1113-1118.
- GEUSKEN R.B.M., LUTEIJN J.M., SCHOOONHOVEN L.M., 1983. Antifeedant activity of some ajugarin derivatives in three lepidopterous species. Experientia 39, 303-404.
- HERNÁNDEZ A., RODRÍGUEZ B., 1981. Cumarinas de *Cachrys trifida* Miller. An Quim 77 (C), 359-360. [In Spanish].
- HERNÁNDEZ-PÉREZ M., RABANAL R.M., DE LA TORRE M.C., RODRÍGUEZ B., 1995. Analgesic, anti-inflammatory, antipyretic and haematological effects of aethiopinone, an *o*-naphthoquinone diterpenoid from *Salvia aethiopsis* roots and two hemisynthetic derivatives. Planta Med 61, 505-509.
- HERNÁNDEZ-PÉREZ M., RABANAL R.M., ARIAS A., DE LA TORRE M.C., RODRÍGUEZ B., 1999. Aethiopinone, an antibacterial and cytotoxic agent from *Salvia aethiopsis* roots. Pharmaceutical Biology 37, 17-21.
- HUESO-RODRÍGUEZ J.A., RODRÍGUEZ B., 1989. A new and efficient route to optically active drimananes. Synthesis of (+)-winterin, (+)-confertifolin, (+)-isodrimenin, and (+)-bicyclofarnesol. Tetrahedron 45, 1567-1576.
- HUESO-RODRÍGUEZ J.A., JIMENO M.L., RODRÍGUEZ B., SAVONA G., BRUNO M., 1983. Abietane diterpenoids from the root of *Salvia phlomoides*. Phytochemistry 22, 2005-2009.
- HUSSEIN A.A., DE LA TORRE M.C., RODRÍGUEZ B., HAMMOUDA F.M., HUSSINEY H.A., 1997. Modified abietane diterpenoids and a methoxylupane derivative from *Salvia palaestina*. Phytochemistry 45, 1663-1668.
- HYUNCHEOL O., HOSUB L., TAEWAN K., KYUNYUN C., HUNTAEG C., TAEOK H., JUNGYANG J., OKSAM J., YOUNCHUL K., YOUNGGAB Y., 2002. Furocoumarins from *Angelica dahurica* with hepatoprotective activity on tacrine-induced cytotoxicity in Hep G2 cells. Planta Med 68, 463-464.
- MARTÍN-PANIZO F., RODRÍGUEZ B., VALVERDE S., 1974. Estructura total revisada del lagascatriol, nuevo diterpeno de *Sideritis angustifolia* Lag. An Quím 70, 164-171. [In Spanish].
- MENDES E., MARCO J.L., RODRÍGUEZ B., JIMENO M.L., LOBO A.M., PRABHAKAR S., 1989. Diterpenoids from *Salvia candelabrum*. Phytochemistry 28, 1685-1690.
- MERRIT A.T., LEY S.V., 1992. Clerodane diterpenoids. Nat Prod Rep 9, 243-287.
- MURILLO R., MUÑOZ D., CABALLERO P., 2006. El potencial de los baculovirus como agentes de control biológico de plagas. Phytoma España 176, 51-63. [In Spanish].
- ORTEGO F., RODRÍGUEZ B., CASTAÑERA P., 1995. Effects of neo-clerodane diterpenes from *Teucrium* on feeding behavior of Colorado potato beetle larvae. J Chem Ecol 21, 1375-1386.
- RAMOS-GUTIÉRREZ J., ORTIZ J.P., VARGAS-OSUNA E., 2004. Susceptibility of first instar *Helicoverpa armigera* (Hübner) and *Earias insulana* (Boisduval) (Lepidoptera: Noctuidae) larvae to *Bacillus thuringiensis* (Berliner) Cry1Ac toxin. Bol San Veg Plagas 30, 239-245. [In Spanish].
- ROBERTSON J.L., PRIESTER H.K., 1992. Pesticide bioassays with arthropods. CRC Press, Boca Raton, USA.
- RODRÍGUEZ B., 1977. 5,4'-Dihydroxy-6,7,8,3'-tetramethoxyflavone from *Sideritis mugronensis*. Phytochemistry 16, 800-801.
- RODRÍGUEZ B., MARTÍN-PANIZO F., 1979. Flavonas de *Sideritis serrata*. An Quím 75(C), 431-432. [In Spanish].
- RODRÍGUEZ M.C., BARLUENGA J., SAVONA G., PIOZZI F., SERVETTAZ O., RODRÍGUEZ B., 1984a. Isoteuflidin, a neo-clerodane diterpenoid from *Teucrium chamaedrys*, and revised structures of teucrins F and G. Phytochemistry 23, 1465-1469.
- RODRÍGUEZ B., FERNÁNDEZ-GADEA F., SAVONA G., 1984b. A rearranged abietane diterpenoid from the root of *Salvia aethiopsis*. Phytochemistry 23, 1805-1806.
- RODRÍGUEZ-HANH L., ESQUIVEL B., CARDENAS J., 1994. Clerodane diterpenoids in Labiateae. In: Progress in the chemistry of organic natural products 63 (Hertz W., Kirby G.W., Moore R.E., Steglich W., Tamm Ch., eds). Springer-Verlag. pp. 107-196.
- RODRÍGUEZ-RODRÍGUEZ M.D., MORENO R., TÉLLEZ M.M., RODRÍGUEZ-RODRÍGUEZ M.P., LASTRES J., 1996. El cultivo del melón bajo plástico en la provincia de Almería. Caracterización y seguimiento de las principales plagas y enfermedades. Phytoma 80, 12-17. [In Spanish].
- SAVONA G., PIOZZI F., RODRÍGUEZ B., 1978. Hispanolone, a new furanoditerpene. Heterocycles 9, 257-261.
- SAVONA G., PATERNOSTRO M.P., PIOZZI F., RODRÍGUEZ B., 1979. Flavonas de *Teucrium pseudochamaepeplis*. An Quím 75(C), 433-434.
- SAVONA G., BRUNO M., PATERNOSTRO M.P., MARCO J.L., RODRÍGUEZ B., 1982. Salviacoccin, a neo-clerodane diterpenoid from *Salvia coccinea*. Phytochemistry 21, 2563-2566.
- SHARABY A., AMMAR N., 1997. Biological activity of extracts of *Tephrosia nubica* (Boiss) Baker against *Spodoptera littoralis* (Boisd.) and *Agrotis ipsilon* (Hufn.). Journal of Agriculture in the Tropics and Subtropics 98, 143-150.
- SIMMONDS M.S.J., BLANEY W.M., LEY S.V., SAVONA G., BRUNO M., RODRÍGUEZ B., 1989. The antifeedant activity of clerodane diterpenoids from *Teucrium*. Phytochemistry 28, 1069-1071.
- SIMMONDS M.S.J., BLANEY W.M., DELLE MONACHE F., MARINI-BETTOLO G.B., 1990. Insect antifeedant activity associated with compounds isolated from species of *Lonchocarpus* and *Tephrosia*. J Chem Ecol 16, 365-380.
- SIMMONDS M.S.J., BLANEY W.M., ESQUIVEL B., RODRIGUEZ-HANH L., 1996. Effect of clerodane-type diterpenoids from *Salvia* sps. on the feeding behaviour of *Spodoptera littoralis*. Pestic Sci 47, 17-23.
- STEVENSON P.C., SIMMONDS M.S.J., YULE M.A., VEITCH N.C., KITE G.C., IRWIN D., LEGG M., 2003. Insect antifeedant furocoumarins from *Tetradium daniellii*. Phytochemistry 63, 41-43.

TÉLLEZ M.M., MORENO R., RODRÍGUEZ-RODRÍGUEZ M.D., RODRÍGUEZ-RODRÍGUEZ M.P., 1996. Evaluación de las principales plagas y enfermedades que afectan al cultivo del calabacín bajo plástico en la provincia de Almería. *Phytoma* 84, 32-37. [In Spanish].

TORRES-VILA L.M., RODRÍGUEZ-MOLINA M.C., LACASA-PLASENCIA A., BIELZA-LINO P. 2002.

Insecticide resistance of *Helicoverpa armigera* to endosulfan, carbamates and organophosphates: the Spanish case. *Crop Prot* 21, 1003-1013.

WELLSOW J., GRAYER R.J., VEITCH N.C., KOKUBUN T., LELLI R., KITE G.C., SIMMONDS M.S.J., 2006. Insect-antifeedant and antibacterial activity of diterpenoids from species of *Plectranthus*. *Phytochemistry* 67, 1818-1825.